

ON RIGHT COIDEAL SUBALGEBRAS

V.K. KHARCHENKO

ABSTRACT. Let H be a character Hopf algebra. Every right coideal subalgebra that contains the coradical has a PBW-basis which can be extended up to a PBW-basis of H .

1. INTRODUCTION

In the present paper we consider right coideal subalgebras in character Hopf algebras, the Hopf algebras generated by skew-primitive semi-invariants. This class includes quantum enveloping algebras of Kac–Moody algebras, all their generalizations (see, G. Benkart, S.-J. Kang, and D. Melville [1]; M. Costantini, and M. Varagnolo [2]; S.-J. Kang [3]), the bosonizations of quantum symmetric algebras related to diagonal braidings, and so on. We prove the following general statement on the structure of the right coideal subalgebras.

Theorem 1.1. *Let H be a character Hopf algebra. Every right coideal subalgebra that contains all group-like elements has a PBW-basis which can be extended up to a PBW-basis of H .*

This theorem is new for Hopf subalgebras too.

2. PRELIMINARIES

2.1. PBW-generators. Let S be an algebra over a field $\mathbf{k}$ and A is a subspace of S with a basis $\{a_j \mid j \in J\}$. A linearly ordered subset $V \subseteq S$ is said to be a *set of PBW-generators of S over A* if there exists a function $h : V \rightarrow \mathbf{Z}^+ \cup \{\infty\}$, called the *height function*, such that the set of all products

$$(2.1) \quad a_j v_1^{n_1} v_2^{n_2} \cdots v_k^{n_k},$$

where $j \in J$, $v_1 < v_2 < \cdots < v_k \in V$, $n_i < h(v_i)$, $1 \leq i \leq k$ is a basis of S . The value $h(v)$ is referred to as the *height* of v in V . If $A = \mathbf{k}$ is the ground field, then we shall call V simply as a *set of PBW-generators of S* .

Definition 2.1. Let V be a set of PBW-generators of S over a subalgebra A . Suppose that the set of all words in V as a free monoid has its own order $\prec$ (that is, $a \prec b$ implies $cad \prec cbd$ for all words a, b, c, d in V).

1) A *leading word* of $s \in S$ is the maximal word $m = v_1^{n_1} v_2^{n_2} \cdots v_k^{n_k}$ that appears in the decomposition of s in the basis (2.1).

2) A *leading term* of s is the sum am of all terms $\alpha_i a_i m$ that appear in the decomposition of s in the basis (2.1), where m is the leading word of s .

1991 *Mathematics Subject Classification.* Primary 16W30, 16W35; Secondary 17B37.

Key words and phrases. Hopf algebra, coideal subalgebra, PBW-basis.

The author was supported by PAPIIT IN 108306-3, UNAM, México.

3) The order $\prec$ is *compatible with the PBW-decomposition related to V* if the leading word of each product $W = w_1 w_2 \cdots w_m$, $w_i \in V \cup \{a_j\}$ (considered as an element of S) is less than or equal to the word W^t that appears from $w_1 w_2 \cdots w_m$ by deletion of all letters a_j .

2.2. Associated graded algebra. Let Γ be a completely ordered additive (commutative) monoid. A Γ -*filtration* on an algebra S is a map D from S to Γ extended by a symbol $-\infty$ with the properties

1. $D(x) = -\infty \iff x = 0$, $D(1 \cdot \mathbf{k}^*) = 0$,
2. $D(x - y) \leq \max\{D(x), D(y)\}$, $D(xy) \leq D(x) + D(y)$.

The element $D(x)$ is the *degree* of x . We denote by S_γ the subspace of all elements of degree $\leq \gamma$, while by S_γ^- the subspace of all elements of degree $< \gamma$.

Example 2.2. (Filtration by constitution). Suppose that an algebra S is generated over a subalgebra A by a set X . That is, there exists a homomorphism $\xi : A\langle X \rangle \rightarrow S$, where $A\langle X \rangle$ is the free product $A * \mathbf{k}\langle X \rangle$. A *constitution* of a word u in $A \cup X$ is a family of non-negative integers $\{m_x, x \in X\}$ such that u has m_x occurrences of x . Certainly almost all m_x in the constitution are zero. We fix an arbitrary complete order, $<$, on the set X .

Let Γ be the free additive (commutative) monoid generated by X . The monoid Γ is a completely ordered monoid with respect to the following order:

$$(2.2) \quad m_1 x_{i_1} + m_2 x_{i_2} + \dots + m_k x_{i_k} > m'_1 x_{i_1} + m'_2 x_{i_2} + \dots + m'_k x_{i_k}$$

if the first from the left nonzero number in $(m_1 - m'_1, m_2 - m'_2, \dots, m_k - m'_k)$ is positive, where $x_{i_1} > x_{i_2} > \dots > x_{i_k}$ in X . We associate a formal degree $D(u) = \sum_{x \in X} m_x x \in \Gamma$ to a word u in $A \cup X$, where $\{m_x \mid x \in X\}$ is the constitution of u . Respectively, if $f = \sum \alpha_i u_i \in A\langle X \rangle$, $0 \neq \alpha_i \in \mathbf{k}$ then

$$(2.3) \quad D(f) = \max_i \{D(u_i)\}.$$

On S we define a Γ -filtration related to $\xi : A * \mathbf{k}\langle X \rangle \rightarrow S$ as follows:

$$(2.4) \quad D(s) = \min\{D(f) \mid \xi(f) = s\}.$$

With every Γ -filtered algebra S a Γ -graded algebra is associated in the obvious way. For each $\gamma \in \Gamma$, write $\text{gr}_\gamma S$ for the linear space S_γ / S_γ^- . Then

$$\text{gr } S = \bigoplus_{\gamma \in \Gamma} \text{gr}_\gamma S,$$

with the product defined by

$$(x + S_\gamma^-)(y + S_\delta^-) = xy + S_{\gamma+\delta}^-.$$

If the filtration is as in Example 2.2, then the algebra $\text{gr } S$ is generated by $\text{gr } X \cong X$ and $\text{gr } A = A$.

Lemma 2.3. *If the associated graded algebra $\text{gr } S$ for a Γ -filtered algebra S has a set of homogeneous PBW-generators, $V = \{v_i\}$, over a subspace $A \subseteq S_0$, then each set of representatives $\hat{v}_i$, $v_i = \hat{v}_i + S_{D(v_i)}^-$, is a set of PBW-generators of S over A .*

This lemma has been proved by S. Ufer [7, Proposition 46] for the case $\Gamma = \mathbf{Z}^+$. In the general case the proof is quite similar.

Lemma 2.4. *Let $\mathbf{U}$ be a subalgebra of a Γ -filtered algebra S . If $\text{gr } S$ is a free left (right) Γ -graded module over $\text{gr } \mathbf{U}$ then S is a free left (right) module over $\mathbf{U}$.*

This is part of the folklore.

2.3. Thin elements and replacement of basis. Suppose that an algebra S has a set V of PBW-generators over a subalgebra A , and $1 \in \{a_j\}$. Let $\prec$ be any complete order of the free monoid of words in V which is compatible with the PBW-decomposition related to V . An element $c \in S$ is said to be a *thin element* if its decomposition in the basis (2.1) has the form

$$(2.5) \quad c = v^m + \sum \alpha_{ij} a_j W_i,$$

where $W_i \prec v^m$, and either m divides the height of v in V , or $h(v) = \infty$.

Let $T \subseteq S$ be a some set of thin elements. Suppose that for each $v \in V$ there exists at most one element $c_v \in T$ with the leading term of the form v^m , $m \geq 1$. One may construct a new set of PBW-generators, P_T , related to T in the following way.

If in T there does not exist an element with the leading term of the form v^m , $m \geq 1$ we include v in P_T and define the height $h_T(v)$ related to T to be equal to $h(v)$.

If there exists an element $c_v \in T$ with the leading term v^m and $m > 1$, we include in P_T both elements: v and c_v . In this case we define the height of v in P_T to be equal to m , while the height of c_v related to P_T is the quotient $h(v)/m$.

If there exists an element $c_v \in T$ with the leading term v (that is, if $m = 1$) then we include in P_T just c_v (that accidentally may be equal to v). In this case the height of c_v with respect to T by definition equals $h(v)$.

We extend on P_T the order $<$ in the natural way: $c_v < w$ if and only if $(v \leq w \text{ \& } c_v \neq w)$; and $c_v < c_w$ if and only if $v < w$. In particular $c_v < v$, provided that $m > 1$, where m is defined by c_v in (2.5).

Lemma 2.5. *The set P_T is a set of PBW-generators of S over A .*

Proof. We have to prove that the monotonous restricted words (2.1) with $v_i \in P_T$ are linearly independent in S , and they span S .

By definition if $\theta_1 < \theta_2 \in P_T$, $\theta_1 = v_1^{m_1} + \dots$, $\theta_2 = v_2^{m_2} + \dots$, then either $v_1 < v_2$, or $v_1 = v_2 = v$ with $\theta_1 = c_v = v^m + \dots$, $\theta_2 = v$. Therefore each monotonous word in P_T has a form

$$(2.6) \quad \theta_1^{n_1} v_1^{r_1} \theta_2^{n_2} v_2^{r_2} \dots \theta_k^{n_k} v_k^{r_k},$$

where $v_1 < v_2 < \dots < v_k$, $\theta_i = c_{v_i} = v_i^{m_i} + \dots$. Of course if $m_i = 1$, then $r_i = 0$, while if $m_i > 1$, then $r_i \geq 0$. The word (2.6) is restricted if and only if $n_i < h_T(\theta_i) = h(v_i)$ in the case $m_i = 1$, and $n_i < h_T(\theta_i) = h(v_i)/m_i$, $r_i < m_i$ otherwise. If we replace $\theta_i := v_i^{m_i} + \dots$ in (2.6) and then develop multiplication, we get a linear combination $W + \sum_q \alpha_q W_q$ of words in $V \cup \{a_j\}$, where

$$(2.7) \quad W = v_1^{n_1 m_1 + r_1} v_2^{n_2 m_2 + r_2} \dots v_k^{n_k m_k + r_k}.$$

Let us as above denote by W^t a word in V that appears from W by deletion of all a_j . Since $\prec$ is a monoidal order, we have $W_q^t \prec W$, for all q . If $m_i > 1$ then

$$n_i m_i + r_i \leq \left(\frac{h(v_i)}{m_i} - 1 \right) m_i + (m_i - 1) = h(v_i) - 1.$$

If $m_i = 1$, again $n_i m_i + r_i = n_i < h(v_i)$. Hence (2.7) is a monotonous restricted word, provided that so is (2.6). Since the order $\prec$ is compatible with the PBW-decomposition related to V , all words $v_1 v_2 \cdots v_k$ that appears in the PBW-decomposition (2.1) of $\sum_q \alpha_q W_q$ are less than (2.7). It is important to note that in this way to different monotonous restricted words in P_T correspond different monotonous restricted words in V .

Suppose that a linear combination, $\Xi = \sum \alpha_{ij} a_j U_i$, over A of monotonous restricted words in P_T equals zero in S . Let us first in Ξ replace each $\theta \in P_T$ by its representation, $\theta := v^m + \cdots$, in terms of PBW-generators V , and next develop the multiplication. We get a linear combination $\Xi' = \sum \beta_k W_k$ of words in $P_T \cup \{a_j \mid j \in J\}$. Since $\prec$ is a monoidal order ($a \prec b$ implies $cad \prec cbd$), the maximal word W_0 among W_k^t appears in the decomposition of summands $a_j U$ with the only word U . If U has form (2.6), then W_0 has form (2.7). Since the order $\prec$ is compatible with the PBW-decomposition related to V , the maximal word of Ξ' in the PBW-decomposition appears just in summands $\alpha_{sj} a_j W_0$, where the equality $U_s = U$ defines the index s . Hence $\Xi' \neq 0$ in S . A contradiction.

To see that monotonous restricted words in $P_T \cup \{a_j\}$ span S we may use a standard induction on words ordered by $\prec$. Indeed, for any monotonous restricted word in V ,

$$(2.8) \quad W = v_1^{s_1} v_2^{s_2} \cdots v_k^{s_k},$$

we have $s_i = n_i m_i + r_i$, $r_i < m_i$, $n_i < h_T(\theta_i)$, where either $\theta_i = c_{v_i} = v_i^{m_i} + \cdots \in T$, or the set T has no elements with the leading word v_i^m , $m \geq 1$ (and hence $\theta_i = v_i \in P_T$, $m_i = 1$). Suppose by induction that values of all super-words smaller then W belong to the linear space span by monotonous restricted words in $P_T \cup \{a_j\}$. The difference between W and (2.6) is a linear combination of words U in $P_T \cup \{a_j\}$ such that U^t are less than W . Due to the comparability of $\prec$ with the PBW-decomposition related to V , the PBW-decomposition of any such a word has only summands $\alpha a_j W'$ with $W' \prec W$. By induction the lemma is proved. $\square$

Remark. In the proof we have seen that the leading term of (2.6) in basis (2.1) equals (2.7).

3. PBW-BASIS OF A CHARACTER HOPF ALGEBRA

Recall that a Hopf algebra H is referred to as a *character* Hopf algebra if the group G of all grouplike elements is commutative and H is generated over $\mathbf{k}[G]$ by skew primitive semi-invariants s_i , $i \in I$:

$$(3.1) \quad \Delta(s_i) = s_i \otimes 1 + g_{s_i} \otimes s_i, \quad g^{-1} s_i g = \chi^{s_i}(g) s_i, \quad g, g_{s_i} \in G,$$

where χ^{s_i} , $i \in I$ are characters of the group G . Let us associate a variable x_i to s_i . For each word u in $X = \{x_i \mid i \in I\}$ we denote by g_u an element of G that appears from u by replacing each x_i with g_{s_i} . In the same way we denote by χ^u a character that appears from u by replacing of each x_i with χ^{s_i} . We define a bilinear skew commutator by the formula

$$(3.2) \quad [u, v] = uv - p_{uv} vu,$$

where by definition $p_{uv} = \chi^u(g_v) = p(u, v)$.

The group G acts on the free algebra $\mathbf{k}\langle X \rangle$ by $g^{-1} u g = \chi^u(g) u$, where u is an arbitrary monomial in X . The skew group algebra $G\langle X \rangle$ has the natural Hopf

algebra structure

$$\Delta(x_i) = x_i \otimes 1 + g_{s_i} \otimes x_i, \quad i \in I, \quad \Delta(g) = g \otimes g, \quad g \in G.$$

The construction of the PBW-basis given in [4] requires I to be finite. In order to get a PBW-basis in the general case one may slightly modify this construction as follows.

We fix a Hopf algebra homomorphism $\xi : G\langle X \rangle \rightarrow H$, $\xi(x_i) = s_i$, $\xi(g) = g$, $i \in I$, $g \in G$, and consider the filtration D related to ξ as defined in Example 2.2. This filtration is compatible with the Hopf algebra structure, that is

$$(3.3) \quad \Delta(H_\gamma) \subseteq \sum_{\delta+\varepsilon=\gamma} H_\delta \otimes H_\varepsilon, \quad \sigma(H_\gamma) \subseteq H_\gamma,$$

where σ is the antipode. Therefore $\text{gr } H$ is also a character Hopf algebra generated by $\mathbf{k}[G]$ and $\xi(X)$.

By Lemmas 2.3 and 2.4 in what follows we may suppose that H is a Γ -graded character Hopf algebra (or, in the other words, it is homogeneous in each of the generators s_i).

On the set of all words in X we fix the lexicographical order with the priority from the left to the right, where a proper beginning of a word is considered to be greater than the word itself. This order is not monoidal: we have $x > x^2$, while $xy < x^2y$ if $y < x$. It has neither ACC, nor DCC: $x > x^2 > \dots > x^n > \dots$; $xy < x^2y < \dots < x^ny < \dots$. By these reasons we need one more order, the *Hall* (or *deg-lex*) one, $u < v$ if $D(u) < D(v)$ or $D(u) = D(v)$ & $u < v$. Since there exist just a finite number of words in X of a given constitution, the Hall order indeed is a complete order of the free monoid.

A non-empty word u is called a *standard* word (or *Lyndon* word, or *Lyndon-Shirshov* word) if $vw > wv$ for each decomposition $u = vw$ with non-empty v, w . A *nonassociative* word is a word where brackets $[,]$ somehow arranged to show how multiplication applies. If $[u]$ denotes a nonassociative word then by u we denote an associative word obtained from $[u]$ by removing the brackets (of course $[u]$ is not uniquely defined by u in general).

The set of *standard nonassociative* words is the biggest set SL that contains all variables x_i and satisfies the following properties.

- 1) If $[u] = [[v][w]] \in SL$ then $[v], [w] \in SL$, and $v > w$ are standard.
- 2) If $[u] = [[[v_1][v_2]][w]] \in SL$ then $v_2 \leq w$.

Every standard word has the only alignment of brackets such that the appeared nonassociative word is standard (the Shirshov theorem). In order to find this alignment one may use the following procedure: The factors v, w of the nonassociative decomposition $[u] = [[v][w]]$ are the standard words such that $u = vw$ and v has the minimal length ([6], see also [5]).

Definition 3.1. A *super-letter* is a polynomial that equals a nonassociative standard word where the brackets mean (3.2). A *super-word* is a word in super-letters.

By Shirshov theorem every standard word u defines the only super-letter, in what follows we will denote it by $[u]$. The order on the super-letters is defined in the natural way: $[u] > [v] \iff u > v$. We should stress that this order is not complete: If $x > y$, there are infinite chains of super-letters

$$[xy] > [xy^2] > \dots > [xy^n] > \dots; [xy] < [x^2y] < \dots < [x^ny] < \dots$$

Nevertheless the Hall order on super-words is still a complete monoidal order.

Definition 3.2. A super-letter $[u]$ is called *hard in H* provided that its value in H is not a linear combination of values of super-words in smaller than $[u]$ super-letters.

Definition 3.3. We say that a *height* of a hard in H super-letter $[u]$ equals $h = h([u])$ if h is the smallest number such that: first, p_{uu} is a primitive t -th root of unity and either $h = t$ or $h = tl^r$, where $l = \text{char}(\mathbf{k})$; and then the value in H of $[u]^h$ is a linear combination of super-words in less than $[u]$ super-letters. If there exists no such number then the height equals infinity.

Theorem 3.4. *Values of all hard in H super-letters form a set of PBW-generators of H over $\mathbf{k}[G]$.*

Proof. If X is a finite set, this statement has been proved in [4, Theorem 2], see a footnote on page 267. In the general case the algebra $G\langle X \rangle$ has a covering by $G\langle X_\alpha \rangle$, where X_α runs through all finite subsets of X . We shall consider each of X_α as an ordered subset of X , and $G\langle X_\alpha \rangle$ as a Γ -graded subalgebra of $G\langle X \rangle$. Respectively H has a covering by the finitely generated Γ -graded character Hopf subalgebras $H_\alpha \stackrel{\text{df}}{=} \xi(G\langle X_\alpha \rangle)$.

By Definition 3.2 every hard in H super-letter $[u]$ is hard in H_α as soon as $u \in G\langle X_\alpha \rangle$. If a super-letter $[v] \in G\langle X_\alpha \rangle$ is not hard in H , then by definition $\xi([v]) = \xi(A)$, where A is a linear combination of super-words in less than $[v]$ super-letters. Since H is Γ -graded, all super-words of the sum A have the same constitution as the word v does. In particular all super-words of A belong to $G\langle X_\alpha \rangle$. Hence $[v]$ is not hard in H_α .

In the same way the height function is independent of α . $\square$

The following statements provide important properties of PBW-basis defined by the hard super-letters.

Lemma 3.5. ([4, Lemma 8]). *The coproduct of a super-letter $[w]$ has a representation*

$$(3.4) \quad \Delta([w]) = [w] \otimes 1 + g_w \otimes [w] + \sum_i \alpha_i g(W_i'') W_i' \otimes W_i'',$$

where W_i' are non-empty words in less than $[w]$ super-letters.

Lemma 3.6. *The Hall order of the super-words is compatible with the PBW-decomposition related to the hard super-letters (see Definition 2.1).*

Proof. Let W be a super-word. There exists the following natural diminishing process of the decomposition. First, in $\mathbf{k}\langle X \rangle$ according to [4, Lemma 7] we decompose the super-word W in a linear combination of smaller monotonous super-words, then, we replace each non hard super-letter with the decomposition of its value in H that exists by Definition 3.2, and again decompose the appeared super-words in linear combinations of smaller monotonous super-words, and so on, until we get a linear combination of monotonous super-words in hard super-letters. If they are not restricted, we may apply Definition 3.3 and repeat the process until we get only monotonous restricted words in hard super-letters. $\square$

Lemma 3.7. *If $[w]$ is a super-letter then*

$$(3.5) \quad \Delta(\xi([w]^m)) = \sum_{j=0}^m \begin{bmatrix} m \\ j \end{bmatrix}_q g_w^{m-j} \xi([w])^j \otimes \xi([w])^{m-j} + \sum_i \alpha_i g(V_i) \xi(U_i) \otimes \xi(V_i),$$

where $\begin{bmatrix} m \\ j \end{bmatrix}_q$ are the q -binomial coefficients with $q = p(w, w)$, and U_i are basis super-words such that the first from the left super-letter is less than $[w]$.

Proof. The m -th power of the right hand side of (3.4) after developing of the product takes the form (3.5), where each of U_i is a product of m super-words some of whom equal to $[w]$ (but not all of them!) and others equal to W'_i 's. Hence U_i as a super-word is less than $[w]^m$ with respect to the lexicographical ordering of words in super-letters. Let us decompose $\xi(U_i)$ in the PBW-basis defined by the hard super-letters. According to Lemma 3.6 we turn to the formula (3.5) where still $U_i < [w]^m$. This implies the required property since the first super-letter in a basis super-word is always the minimal one. $\square$

4. RIGHT COIDEAL SUBALGEBRAS

Theorem 4.1. *Let H be a character Hopf algebra. Every right coideal subalgebra $\mathbf{U} \supseteq \mathbf{k}[G]$ has a PBW-basis that may be extended up to a PBW-basis of H .*

By Lemma 2.3 it suffices to show that $\text{gr}\mathbf{U}$ as a subalgebra of $\text{gr}H$ has a required basis. Since $\text{gr}\mathbf{U}$ is a right coideal subalgebra of $\text{gr}H$, we still may suppose that H is a Γ -graded character Hopf algebra and $\mathbf{U}$ is a Γ -graded coideal subalgebra.

Lemma 4.2. *Let $w \in \mathbf{U}$. If $\pi, \mu : H \rightarrow H$ are linear maps such that $(\pi \otimes \mu)\Delta(w) = a \otimes b$ with $b \neq 0$, then $a = \pi(c)$ for some $c \in \mathbf{U}$.*

Proof. We have $(\pi \otimes \mu)\Delta(\mathbf{U}) \subseteq \pi(\mathbf{U}) \otimes H$. Hence $a \otimes b \in \pi(\mathbf{U}) \otimes H$, and $a \in \pi(\mathbf{U})$. $\square$

We shall use this evident statement as a basic tool. Note that once we have a PBW-basis of H , to define a linear map it suffices to fix its values on the restricted monotonous words in an arbitrary way.

Let $[u]$ be a hard super-letter. Suppose that in $\mathbf{U}$ there exists an element c with the leading super-word $[u]^m$, $m \geq 1$. Since $G \subseteq \mathbf{U}$, we may suppose that the super-word $[u]^m$ appears one time with the trivial coefficient:

$$(4.1) \quad c = \xi([u]^m + \sum_j \alpha_j g_j [u]^m + \sum_i \alpha_i g_i W_i R_i),$$

where W_i are nonempty basis super-words in less than $[u]$ super-letters, while R_i are basis super-words in greater than or equal to $[u]$ super-letters, $\alpha_i, \alpha_j \in \mathbf{k}, g_i, g_j \in G, g_j \neq 1$.

Denote by ι the natural projection $H \rightarrow \mathbf{k}$, $\iota(g\xi(W)) = 0$, unless $g = 1, W = \emptyset$. Since $\Delta(gW) = (g \otimes g)\Delta(W)$, we have

$$(\text{id} \otimes \iota)(\Delta(c)) = \xi([u]^m + \sum_{g_i=1} \alpha_i W_i R_i) \otimes 1.$$

Thus, by Lemma 4.2

$$(4.2) \quad c' = \xi([u]^m + \sum_{g_i=1} \alpha_i W_i R_i) \in \mathbf{U}.$$

In what follows we fix the notation c_u for one of the elements from $\mathbf{U}$ of the form (4.2) that has the minimal possible m .

Lemma 4.3. *In the representation (4.2) of the chosen element c_u either $m = 1$, or p_{uu} is a primitive t -th root of unity and $m = t$ or (in the case of positive characteristic) $m = t(\text{char } \mathbf{k})^s$. In particular c_u is a thin element, see (2.5).*

Proof. If $m = 1$ there is nothing to prove. Let $m > 1$. For each $k, 1 \leq k < m$ we consider the following linear map set up on the PBW-basis of super-words.

$$(4.3) \quad \pi_k(g\xi(W)) = \begin{cases} 0, & \text{if } W \prec [u]^k, \\ g\xi(W), & \text{otherwise;} \end{cases}$$

By means of formula (3.5) we have

$$(4.4) \quad (\pi_k \otimes \pi_{m-k})\Delta(\xi([u]^m)) = \left[\begin{matrix} m \\ k \end{matrix} \right]_q g_u^{m-k} \xi([u]^k) \otimes \xi([u]^{m-k}).$$

By Lemma 3.5 and Lemma 3.6 the coproduct $\Delta(\xi(W_i))$ is the sum of tensors $g\xi(W'_i) \otimes \xi(W''_i)$, where all W'_i are the basis super-words lexicographically smaller than $[u]^m$ with the only exception that equals $g(W_i) \otimes \xi(W_i)$. Hence $(\pi_k \otimes \pi_{m-k})\Delta(\xi(W_i R_i))$ is a sum of tensors of the form $g\pi_k(\xi(R'_i)) \otimes \pi_{m-k}(\xi(W_i R''_i))$. Again by Lemma 3.6 we have $\pi_{m-k}(\xi(W_i R''_i)) = 0$. Hence

$$(4.5) \quad (\pi_k \otimes \pi_{m-k})\Delta(\xi(W_i R_i)) = 0.$$

Thus we may write

$$(4.6) \quad (\pi_k \otimes \pi_{m-k})\Delta(c_u) = \left[\begin{matrix} m \\ k \end{matrix} \right]_q g_u^{m-k} \xi([u]^k) \otimes \xi([u]^{m-k}).$$

If $\left[\begin{matrix} m \\ k \end{matrix} \right]_q \neq 0$, then by Lemma 4.2 we find $c \in \mathbf{U}$ such that $\pi_k(c) = g_u^{m-k} \xi([u]^k)$. By definition of π_k this equality means that the PBW-decomposition of $g_u^{k-m} c$ has the form (4.2) with k in place of m . Since by the choice of c_u the number m is minimal, we get $\left[\begin{matrix} m \\ k \end{matrix} \right]_q = 0, 1 \leq k < m$. This system of equations implies $q^m = 1$ and either m equals the multiplicative order t of q or $m = t(\text{char } \mathbf{k})^s$. $\square$

By Lemma 2.5 the set T of all above defined elements c_u has an extension up to a set P_T of PBW-generators of H over $\mathbf{k}[G]$. Now Theorem 4.1 follows from the proposition below.

Proposition 4.4. *An element $c \in H$ belongs to $\mathbf{U}$ if and only if all PBW-generators in the PBW decomposition of c with respect to P_T belong to T . In particular T is a set of PBW-generators of $\mathbf{U}$ over $\mathbf{k}[G]$.*

To prove this statement we will need some additional properties of the PBW-basis defined by P_T . We extend the order “ $<$ ” already defined on P_T onto the set of all words in P_T as the lexicographical order. The order “ $\prec$ ” is the Hall order induced by the degree function D ; that is, $W \prec U$ if and only if either $D(W) < D(U)$, or $D(W) = D(U)$ & $W < U$. We have to stress that the order on the set of letters P_T differs from the order on the set of one-letter words P_T .

We start with connections between these two PBW-decompositions.

Lemma 4.5. *The leading term of a monotonous restricted word in P_T ,*

$$(4.7) \quad \theta_1^{n_1} [u_1]^{r_1} \theta_2^{n_2} [u_2]^{r_2} \cdots \theta_k^{n_k} [u_k]^{r_k},$$

under the PBW-decomposition related to the hard super-letters equals

$$(4.8) \quad [u_1]^{n_1 m_1 + r_1} [u_2]^{n_1 m_2 + r_2} \dots [u_k]^{n_k m_k + r_k},$$

where $\theta_i = [u_i]^{m_i} + \dots$. Conversely if

$$(4.9) \quad W = [u_1]^{s_1} [u_2]^{s_2} \dots [u_k]^{s_k}$$

is a monotonous restricted super-word in hard super-letters, then its leading term in the decomposition with respect to P_T equals (4.7), where $n_i = \lfloor s_i / m_i \rfloor$, $r_i = s_i - n_i m_i$.

Proof. The first part of the lemma has been proved in Lemma 2.5. The second part follows by induction on super-words ordered by the Hall order. Indeed the difference E between W and (4.7) is a linear combination of super-words that are less than W . By Lemma 3.6 all basis super-words in the PBW-decomposition of E are less than W . It remains to note that if $W' \prec W$ is another basis super-word, then the word (4.7) related to W' is less than that related to W . $\square$

Lemma 4.6. *The Hall order on the words in P_T is compatible with the PBW-decomposition related to P_T (see Definition 2.1).*

Proof. Let W be a word in P_T . The word W has the form (4.7), where $\theta_i = c_{u_i} = \xi([u_i]^{m_i} + \dots)$, while u_i 's are not necessary increase from the left to the right. If we replace $\theta_i := [u_i]^{m_i} + \dots$ in (4.7) and then develop the multiplication in $G\langle X \rangle$, we get a linear combination Σ over $\mathbf{k}[G]$ of super-words with the leading term

$$(4.10) \quad [u_1]^{n_1 m_1 + r_1} [u_2]^{n_1 m_2 + r_2} \dots [u_k]^{n_k m_k + r_k}.$$

By Lemma 3.6 each super-word W_1 in the PBW-decomposition of $\xi(\Sigma)$ is less than or equal to (4.10). That is, if W_1 is different from (4.10), we may write

$$(4.11) \quad W_1 = [u_1]^{n_1 m_1 + r_1} [u_2]^{n_2 m_2 + r_2} \dots [u_s]^{n_s m_s + r_s} [u_{s+1}]^t [v] \dots,$$

here $[v]$ is the first from the left super-letter where W_1 differ from (4.10), and $0 \leq t < n_{s+1} m_{s+1} + r_{s+1}$. Since W_1 is monotonous, we have $v > u_{s+1}$, provided that $t \neq 0$. However in this case W_1 is greater than (4.10). Hence $t = 0$, and $u_{s+1} > v \geq u_s \geq u_{s-1} \geq \dots \geq u_1$.

If all inequalities among u_i 's in the above chain are strict, then according to Lemma 4.5, all words of the PBW-decomposition with respect to P_T of $\xi(W_1)$ are less than or equal to

$$(4.12) \quad \theta_1^{n_1} [u_1]^{r_1} \theta_2^{n_2} [u_2]^{r_2} \dots \theta_s^{n_s} [u_s]^{r_s} \theta_v,$$

where $\theta_v = c_v$, or $\theta_v = [v]$. Hence they are less than W .

If not all inequalities are strict, say $u_1 < u_2 < \dots < u_p = \dots = u_q < u_{q+1}$, $q \leq s$, then again by Lemma 4.5 the P_T -leading word of $\xi(W_1)$ starts with

$$(4.13) \quad \theta_1^{n_1} [u_1]^{r_1} \dots \theta_{p-1}^{n_{p-1}} [u_{p-1}]^{r_{p-1}} \theta_p^{(n_p + \dots + n_q + n_0) m_p} [u_p]^{r_0},$$

where $r_p + \dots + r_q = n_0 m_p + r_0$, $0 \leq r_0 < m_p$. It is still less than W since by definition $\theta_p < [u_p]$, provided that $m_p > 1$. $\square$

Lemma 4.7. *The coproduct of each $\theta \in P_T$ has a representation*

$$(4.14) \quad \Delta(\theta) = \theta \otimes 1 + g_\theta \otimes \theta + \sum_i \alpha_i g_i W'_i \otimes W''_i,$$

where W'_i, W''_i are restricted monotonous words (products) in P_T , and for every i either W'_i or W''_i starts with a letter that is less than θ .

Proof. Let $\theta = \xi([u]^m + \sum_i W_i R_i)$ be the decomposition of $\theta \in P_T$ with respect to the hard super-letters. By Lemma 3.7 and Lemma 4.3, we have

$$\Delta(\xi([u]^m)) - \xi([u]^m) \otimes 1 - g_u^m \otimes \xi([u]^m) = \sum_i \alpha_i g_i \xi(U_i) \otimes \xi(V_i),$$

where U_i are basis super-words starting with smaller than $[u]$ super-letters. By the second part of Lemma 4.5 each summand of the decomposition of $\xi(U_i)$ in the PBW-basis defined by P_T starts by c_w or $[w]$ with $w < u$.

Let $W_i = [w]U$, where $[w] < [u]$. By Lemma 3.5 we have

$$\Delta(W_i R_i) = \sum_j (V'_{ij} \otimes V''_{ij}) \Delta(U R_i) + (g_w \otimes [w]) \Delta(U R_i),$$

where all V'_{ij} are non empty super-words in less than $[u]$ super-letters. All left hand sides of the tensors $(V'_{ij} \otimes V''_{ij}) \Delta(U R_i)$ start with smaller than $[u]$ super-letters. Hence by Lemma 3.6 it remains to use the converse part of Lemma 4.5.

In perfect analogy, if $\Delta(U R_i) = \sum_j U'_{ij} \otimes U''_{ij}$, then $(g_w \otimes [w]) \Delta(U R_i) = g_w U'_{ij} \otimes [w] U''_{ij} R_i$. Therefore the right hand side of each tensor that appears in the P_T -decomposition of $(g_w \otimes \xi([w])) \Delta(\xi(U R_i))$ starts with a letter that is less than θ . $\square$

Lemma 4.8. *Let $\theta \in P_T$. The coproduct of θ^n has a decomposition*

$$(4.15) \quad \Delta(\theta^n) = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_q g_\theta^{n-j} \theta^j \otimes \theta^{n-j} + \sum_i \alpha_i g_i W'_i \otimes W''_i,$$

where $\begin{bmatrix} n \\ j \end{bmatrix}_q$ are q -binomial coefficients with $q = p(\theta, \theta)$, W'_i, W''_i are restricted monotonous words in P_T , and for every i either W'_i or W''_i starts with a letter that is less than θ .

Proof. If we develop multiplication in the n -th power of the right hand side of (4.14), then we get the first sum of (4.15) and a $\mathbf{k}[G]$ -linear combination of tensors $\theta^i W' \otimes \theta^j W''$, where i and j may be zero, but either W' or W'' starts with a less than θ letter. Let it be W' . By Lemma 4.6 the PBW-decomposition of $\theta^i W'$ has only words that are less than $\theta^i W'$ in lexicographical order. Since these words are monotonous, no one of them may start with a letter that is greater than or equal to θ . The lemma is proved. $\square$

Proof of Proposition 4.4. Let in contrary $c \in \mathbf{U}$ be an element of the minimal degree whose decomposition in P_T -basis has super-letters $[u] \in P_T \setminus T$. Since $\mathbf{U}$ is a subalgebra, we may suppose that each term of the decomposition has such a super-letter.

Let $U = \theta_1^{n_1} \theta_2^{n_2} \dots \theta_k^{n_k}$, $\theta_1 < \theta_2 < \dots < \theta_k$ be the leading word of c in the P_T -basis. Since $G \subseteq \mathbf{U}$, we may suppose that U appears one time with the trivial coefficient:

$$(4.16) \quad c = U + \sum_j \alpha_j g_j U + \sum_i \beta_i g_i U_i,$$

where $D(U_i) = D(U)$, $\alpha_j, \beta_i \in \mathbf{k}$, $g_i, g_j \in G$, $g_j \neq 1$.

Denote by ι the natural projection $H \rightarrow \mathbf{k}$, $\iota(gW) = 0$, unless $g = 1$, $W = \emptyset$. Here W is an arbitrary restricted monotonous word in P_T . Since $\Delta(gW) = (g \otimes g) \Delta(W)$,

we have

$$(\text{id} \otimes \iota)(\Delta(c)) = (U + \sum_{g_i=1} \beta_i U_i) \otimes 1,$$

Thus, by Lemma 4.2

$$(4.17) \quad c' = U + \sum_{g_i=1} \beta_i U_i \in \mathbf{U}$$

where $U_i < U$. To get a contradiction we consider two cases.

Case 1. Suppose that $\theta_k \notin T$; that is, $\theta_k = [u]$ and in $\mathbf{U}$ there does not exist an element with the leading super-word $[u]^m$, $1 \leq m \leq n_k$. Let us define the following two linear maps set up on the PBW-basis related to P_T .

$$(4.18) \quad \pi(gW) = \begin{cases} 0, & \text{if } W \prec \theta_k^{n_k}; \\ gW, & \text{otherwise.} \end{cases}$$

$$(4.19) \quad \nu(gW) = \begin{cases} 0, & \text{if } W \prec \theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}; \\ gW, & \text{otherwise.} \end{cases}$$

Let us show first that

$$(4.20) \quad (\pi \otimes \nu)(\Delta(U)) = g\theta_k^{n_k} \otimes \theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}, \quad g \in G.$$

Since the word U is monotonous, by Lemma 4.8 we have

$$(4.21) \quad \Delta(\theta_i^{n_i}) = \sum_{j_i} \alpha_{j_i} g_{j_i} W'_{j_i} \otimes W''_{j_i}, \quad i < k,$$

where for each j_i , $i \leq k$ either W'_{j_i} starts by a letter from P_T that is less than θ_k , or W''_{j_i} starts by a letter from P_T that is less than θ_i , with the only exception $W'_{j_i} = \emptyset$, $W''_{j_i} = \theta_i^{n_i}$. In the same way

$$(4.22) \quad \Delta(\theta_k^{n_k}) = \sum_{j_k} \alpha_{j_k} g_{j_k} W'_{j_k} \otimes W''_{j_k},$$

where either W'_{j_k} or W''_{j_k} starts by a letter from P_T that is less than θ_k , with exceptions of the form $W'_{j_k} = \theta_k^j$, $W''_{j_k} = \theta_k^{n_k-j}$.

$\Delta(U)$ is a right linear combination over $\mathbf{k}[G]$ of tensors

$$(4.23) \quad E = W'_{j_1} W'_{j_2} \cdots W'_{j_k} \otimes W''_{j_1} W''_{j_2} \cdots W''_{j_k}.$$

Let W'_{j_s} be the first from the left nonempty factor in E .

If, first, $s \neq k$, then the tensor get the form

$$E = W'_{j_s} \cdots W'_{j_k} \otimes \theta_1^{n_1} \cdots \theta_{s-1}^{n_{s-1}} W''_{j_s} \cdots W''_{j_k}.$$

If W'_{j_s} starts by a less than θ_k letter, then the left hand side of E is less than $\theta_k^{n_k}$, hence by Lemma 4.6 we get $(\pi \otimes \nu)(E) = 0$.

If W''_{j_s} starts by a less than θ_s letter, then the right hand side is less than $\theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}$, and again $(\pi \otimes \nu)(E) = 0$.

If, next, $s = k$, then the tensor has the form

$$E = W'_{j_k} \otimes \theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}} W''_{j_k}.$$

Since $D(W'_{j_k}) = n_k D(\theta_k) - D(W''_{j_k}) < n_k D(\theta_k)$, unless $W''_{j_k} = \emptyset$, we get $\pi(W'_{j_k}) = 0$ with the only exception $W'_{j_k} = \theta_k^{n_k}$. Thus $\pi \otimes \nu$ kills all tensors in the PBW-decomposition of $\Delta(U)$ except one of them. This proves (4.20).

Let us show, further, that

$$(4.24) \quad (\pi \otimes \nu)(U_i) = 0.$$

Since $U_i < U$, we have $U_i = \theta_1^{n_1} \cdots \theta_s^{n_s} \theta_{s+1}^r \eta \cdots$, where $0 \leq r < n_{s+1}$, $\eta < \theta_{s+1}$. If $r \neq 0$, then $\theta_{s+1} < \eta$ since the word U_i is monotonous. Thus $r_s = 0$; that is,

$$U_i = \theta_1^{n_1} \cdots \theta_s^{n_s} \eta \cdots, \quad \eta < \theta_{s+1}.$$

By means of Lemma 4.7 we have

$$(4.25) \quad \Delta(\eta) = \sum_j \alpha_j g_j V_j' \otimes V_j'',$$

where for each j , either V_j' or V_j'' starts by a letter from P_T which is less than θ_{s+1} . Thus $\Delta(U_i)$ is a right linear combination over $\mathbf{k}[G]$ of tensors

$$(4.26) \quad E = W_{j_1}' \cdots W_{j_s}' V_j' \cdots \otimes W_{j_1}'' \cdots W_{j_s}'' V_j'' \cdots.$$

If $W_{j_1}' = \cdots = W_{j_s}' = V_j' = \emptyset$, then the right hand side takes up the form $\theta_1^{n_1} \cdots \theta_s^{n_s} \eta \cdots$ which is less than $\theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}$, hence $(\pi \otimes \nu)(E) = 0$.

Let W_{j_r}' , $r \leq s$ be the first from the left nonempty word in E . The tensor takes up the form

$$E = W_{j_r}' \cdots W_{j_s}' V_j' \cdots \otimes \theta_1^{n_1} \cdots \theta_{r-1}^{n_{r-1}} W_{j_r}'' \cdots W_{j_s}'' V_j'' \cdots.$$

Since either W_{j_r}' starts with a less than θ_k letter, or W_{j_r}'' starts with a less than θ_r letter, again $(\pi \otimes \nu)(E) = 0$.

If the first from the left nonempty factor is V_j' , then in the same way either V_j' starts with a letter $\leq \eta < \theta_{s+1} \leq \theta_k$, or V_j'' does. In both cases $(\pi \otimes \nu)(E) = 0$.

Thus $(\pi \otimes \nu)(\Delta(U_i)) = 0$. Now taking into account (4.17), we get

$$(4.27) \quad (\pi \otimes \nu)(\Delta(c')) = g \theta_k^{n_k} \otimes \theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}, \quad g \in G.$$

Lemma 4.2 implies that in $\mathbf{U}$ there exists an element c'' such that $\pi(c'') = g \theta_k^{n_k}$. Definition (4.18) of π shows that $\theta_k^{n_k} = [u]^{n_k}$ is the leading word of c'' in the P_T -decomposition. By Lemma 4.5 the leading super-word of c'' equals $[u]^{n_k}$. According to the construction of T , there exists an element $c_u \in T$ of the form $c_u = [u]^m + \cdots$, $m \leq n_k$. This contradicts the conditions of the first case ($[u] = \theta_k \notin T$, $h_T([u]) > n_k$).

Case 2. Suppose that $\theta_k \in T$. In this case $\theta_k \in \mathbf{U}$, therefore one of the letters $\theta_1, \dots, \theta_{k-1}$ does not belong to T . By the inductive supposition no one element with the leading word $\theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}$ belongs to $\mathbf{U}$. At the same time, in perfect analogy with the first case (up to left-right symmetry in the consideration of tensors), we have

$$(\nu \otimes \pi)(\Delta(c')) = g \theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}} \otimes \theta_k^{n_k}.$$

Hence by Lemma 4.2 there exists $c'' \in \mathbf{U}$ such that $\nu(c'') = g \theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}$. Definition (4.19) of ν implies that the leading word of c'' indeed equals $\theta_1^{n_1} \cdots \theta_{k-1}^{n_{k-1}}$. A contradiction. Proposition 4.4 and Theorem 4.1 are completely proved.

REFERENCES

1. G. Benkart, S.-J. Kang, and D. Melville, *Quantized enveloping algebras for Borchers superalgebras*, TAMS, 350, N8(1998), 3297–3319.
2. M. Costantini, M. Varagnolo, Multiparameter quantum function algebra at roots of 1, Math. Ann. 306, N4(1996), 759–780.

3. S.-J. Kang, *Quantum deformations of generalized Kac-Moody algebras and their modules*, Journal of Algebra, 175(1995), 1041–1066.
4. V.K. Kharchenko, *A quantum analog of the Poincaré-Birkhoff-Witt theorem*, Algebra i Logika, 38, N4(1999), 476–507. English translation: Algebra and Logic, 38, N4(1999), 259–276 (QA/0005101).
5. M. Lothaire, *Combinatorics on Words*, Encyclopedia of Mathematics and its Applications, Addison-Wesley Publ. Co., 1983.
6. A.I. Shirshov, *Some algorithmic problems for Lie algebras*, Sibirskii Math. J., 3(2)(1962), 292–296.
7. S. Ufer, *PBW bases for a class of braided Hopf algebras*, Journal of Algebra, 280, N1(2004), 84–119.

UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, FACULTAD DE ESTUDIOS SUPERIORES CUAUTITLÁN, PRIMERO DE MAYO S/N, CAMPO 1, CIT, CUAUTITLÁN IZCALLI, ESTADO DE MÉXICO, 54768, MEXICO

E-mail address: vlad@servidor.unam.mx